



ISSN: 2455-9377

Carbon balance in Maize (*Zea mays* L.) with various combinations of organic and inorganic fertilizers

Jauhari Syamsiyah^{1*}, Ganjar Herdiansyah¹, Lidya Zaela Wijaya², Suntoro¹, Sri Hartati¹

¹Department of Soil Science, Faculty of Agriculture, Sebelas Maret University, Surakarta, Central Java, Indonesia,

²Undergraduate Program, Soil Science, Faculty of Agriculture, Sebelas Maret University, Surakarta, Central Java, Indonesia

ABSTRACT

Fertilization in agricultural cultivation has an important role in the carbon cycle. This study aims to evaluate using organic fertilizers with inorganic fertilizers that can produce high carbon sequestration with low CO₂ emission levels in maize (*Zea mays* L.) cultivation. There were seven combinations of NPK and organic fertilizers, namely 1/4 NPK + 1 OF (C), 1/2 NPK + 1 OF (D), 3/4 NPK + 1 OF (E), 1 NPK + 1 OF (F), 3/4 NPK + 1/4 OF (G), 3/4 NPK + 1/2 OF (H), 3/4 NPK + 3/4 OF (I), and two controls, namely no fertilizer (A) and standard NPK (350 kg/ha, SP36 150 kg/ha, KCl 75 kg/ha) (B). Organic fertilizer was applied one week before planting, SP-36 and KCl fertilizer at planting, and urea fertilizer at 0, 14, and 28 HST. Maize was planted with a spacing of 20 x 70 cm. Parameters observed included CO₂ emissions, soil pH, C-Organic, C-microbial, and plant growth parameters. The combined use of NPK and organic fertilizer significantly increased soil carbon stocks (33.25-64.04 Mg/ha) and carbon sequestration by plants (3.76-5.98 Mg/ha). Therefore, using organic and inorganic fertilizer can be considered more environmentally friendly and effective in managing carbon balance on farmland and has great potential to contribute to climate change mitigation through increased soil and plant carbon sequestration.

KEYWORDS: Carbon Sequestration, CO₂ Emissions, Soil C-Stock

Received: May 23, 2024
Revised: August 11, 2024
Accepted: August 15, 2024
Published: September 07, 2024

*Corresponding author:
Jauhari Syamsiyah
E-mail: ninukts@staff.uns.ac.id

INTRODUCTION

Rising temperatures due to global warming have led to a decline in production on agricultural land (Chauhan *et al.*, 2014). A nearly fifty percent increase in the concentration of carbon dioxide (CO₂) in the atmosphere, from 280 ppm in 1750 to 414.7 ppm in May 2019 (Nema *et al.*, 2012; Wu *et al.*, 2020), often considered to be a significant contributor to global warming and climate change (Li *et al.*, 2013; Al-Ghussain, 2019). One of the causes of increased carbon dioxide (CO₂) concentrations is agricultural activities such as tillage and fertilization.

Fertilization using inorganic fertilizers has contributed to increasing CO₂, this is one of the triggers for global warming in the agricultural sector, as is the use of organic fertilizers. Inorganic fertilizers can contribute to CO₂ emissions through increased root respiration (Chu *et al.*, 2007). At the same time, it can also increase the absorption of CO₂ by the leaves through photosynthesis (Kartikawati & Nursyamsi, 2013). Using

organic fertilizers can also contribute to CO₂ emissions from mineralization and microbial activity (Maswar *et al.*, 2014). However, the use of organic fertilizer has other advantages, namely organic fertilizers can increase soil nutrient availability, microbial activity, and microbial diversity (Yuan *et al.*, 2017), and increase soil C-Stock (Zuo *et al.*, 2023). Although the use of organic fertilizer can increase CO₂ emissions, on the other hand, it can also reduce CO₂ emissions by increasing C-stock. Various studies on CO₂ emissions and C sequestration have been conducted, such as CO₂ emissions in pineapple cultivation in the tropics (Liang *et al.*, 2022), the cultivation of wheat, rice, and maize (Shakoor *et al.*, 2022), in rice-wheat rotation (Zhang *et al.*, 2016), and onion plants (Suwandi *et al.*, 2015).

The increase in the amount of CO₂ must be controlled by increasing the amount of CO₂ absorbed by plants and suppressing the emission of CO₂ back into the air (Ferrini *et al.*, 2020). Thus, it is crucial to quantify the variability of carbon stocks and fluxes through on-farm carbon balance (Béziat *et al.*, 2009; Wang *et al.*, 2015). However, most previous research on

Copyright: © The authors. This article is open access and licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted, use, distribution and reproduction in any medium, or format for any purpose, even commercially provided the work is properly cited. Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

carbon balance has focused on forest and grassland ecosystems and only a tiny proportion of agricultural land (Machado *et al.*, 2016). The use of organic fertilizer is the main choice in efforts to reduce CO₂ emissions. Several previous studies have reported that the use of organic amendments can produce the highest carbon balance in horticultural crops (Persiani *et al.*, 2019), upland rice (Dossou-Yovo *et al.*, 2016), and wheat crops (Roß *et al.*, 2022). In practice, the use of organic fertilizer affects the availability of nutrients in the soil because it is slow-release. In fact, plants need nutrients for growth, so the addition of inorganic fertilizer is still needed. The use of inorganic fertilizers is related to carbon absorption. According to (Rasse *et al.*, 2005; Russell *et al.*, 2009) increasing the nitrogen fertilization dose is also recommended to increase soil carbon sequestration due to increased aboveground biomass. However, the increased carbon uptake resulting from increased nitrogen fertilization may be offset by high CO₂ emissions to the atmosphere (Liu *et al.*, 2019). More effective efforts are needed to reduce CO₂ emissions, one of which is by integrating organic and inorganic fertilizers. Based on these data, it appears that different fertilization patterns have other impacts on carbon balance. However, there is little information on the integrated assessment of the effects of organic and inorganic fertilizer balances on soil C balance processes. This study aims to evaluating the use of organic fertilizers with inorganic fertilizers that can produce high carbon sequestration with low CO₂ emission levels in maize (*Z. mays* L.) cultivation.

MATERIALS AND METHODS

The research was conducted in Klaten, Central Java (7°63'70.053" LS 110°70'04.27" East) from August 2021 to March 2022 on Inceptisol soil with the characteristics presented in Table 2. This field experiment used a randomized complete block design (RCBD) consisting of 7 treatments of NPK and organic fertilizer combinations, namely 1/4 NPK + 1 organic fertilizer (C), 1/2 NPK + 1 organic fertilizer (D), 3/4 NPK + 1 organic fertilizer (E), 1 NPK + 1 organic fertilizer (F), 3/4 NPK + 1/4 organic fertilizer (G), 3/4 NPK + 1/2 organic fertilizer (H), 3/4 NPK + 3/4 organic fertilizer (I) and two control treatments consisting of no fertilizer (A) and standard NPK (350 kg/ha, SP36 150 kg/ha, KCl 75 kg/ha) (B) (Table 1). Crumb organic fertilizer with the properties in Table 3 was applied one week before planting, SP-36, and KCl fertilizers at planting while urea fertilizer at 0, 14, and 28 DAP. The experimental plot size was 20 m² (4 m x 5 m) with a planting distance of 20 cm x 70 cm. Maintenance was carried out as usual by farmers.

CO₂ emissions were measured three times at 35, 49, and 63 HST with a modified Alkali Trap method (Figure 1) (Franzluebbers & Veum, 2020). Measurements were taken in the morning to minimize daily variations and obtain representative daily average soil CO₂ emissions (Xu & Qi, 2001). Samples of disturbed soil were taken at a depth of 0-20 cm to measure various parameters, including soil pH (potentiometry), which was measured three times along with emission sampling (Oladele & Adetunji, 2021), C-Organic (Walkley and Black) (Romadhan *et al.*, 2022), and C-microbe (Fumigation and Extraction) (Rotbart *et al.*, 2017),

while Bulk density is measured using undisturbed soil samples taken with a sampling ring (Gravimetric method) (Irawan *et al.*, 2022). Dry plant biomass was calculated using two plant samples in each plot (Gravimetric Method) (Ekowati & Nasir, 2011).

Soil CO₂ emissions are calculated using the equation:

$$\text{Soil CO}_2 \text{ emissions } \left(\frac{\text{mg}}{\text{m}^2} \right) = (C - S) \times N \times E \quad (\text{Ölinger et al., 1996}) \quad (1)$$

The formula calculates soil C-stock:

$$\text{SOC Stock (Mg C ha}^{-1}\text{)} = [\text{SOC \%}] \times \text{BD 0-20 cm (g/cm}^3\text{)} \times \text{D (0,2 m)} \quad (\text{Li et al., 2017; Abera et al., 2021}) \quad (2)$$

The amount of carbon in plant biomass is calculated using the following formula:

$$C = \text{Biomassa} \times \% \text{ C Biomassa} \quad (\text{IPCC, 2006}) \quad (3)$$

Carbon balance is calculated using the formula:

$$\text{Carbon balance: C soil input - C soil output} \quad (\text{Singh et al., 2009; Sainju et al., 2021}) \quad (4)$$

The research data were analyzed using Analysis of Variance (Anova) with a confidence level of 95%. If the results of Anova achieved are significantly different, it is continued with Duncan's Multiple Range Test (DMRT) to determine differences between treatment means and Pearson -elation test to see the relationship between variables.

RESULTS AND DISCUSSION

Soil CO₂ emission is the release of CO₂ from root respiration, soil microbes, and fauna in the subsoil through biological processes by converting organic carbon into CO₂ (Ebrahimi *et al.*, 2019). CO₂ emissions were significantly affected by organic and inorganic fertilizers ($p=0.000$). The average CO₂ emissions produced ranged from 1.92-6.16 mg CO₂/m²/week and increased with increasing doses of organic fertilizer. This result is consistent with Liu *et al.* (2011) that the balanced application of organic and inorganic fertilizers resulted in higher CO₂ emissions than NPK-only and no-fertilizer treatments due to increased microbial activity from organic amendments. Organic fertilizers applied can increase soil C (Zulkarnain *et al.*, 2013; Syamsiyah *et al.*, 2023a, b) and will supply nutrients for microorganisms as well as stimulate heterotrophic soil respiration and CO₂ emission (Wang *et al.*, 2022). This can be seen in the positive correlation between CO₂ emissions and soil organic C ($r=0.834^{**}$) and C-microbe ($r=0.527^{**}$). The same thing was also stated by De Urzedo *et al.* (2013) and Syamsiyah *et al.* (2019) that CO₂ emissions on farmland increase after organic fertilizer application due to the rise in the supply of labile C to soil microorganisms, thereby increasing the activity of microorganisms and resulting in an increase in soil CO₂ emissions.

Table 1: Experimental treatment design

S. No.	Treatment	Organic Fertilizer (kg/h)	NPK fertilizer (kg/h)		
			Urea	SP36	KCl
1	No fertilizer	0	0	0	0
2	Standard NPK	0	350	150	75
3	¼ NPK+1 Organic Fertilizer	10000	87.5	37.5	18.75
4	½ NPK+1 Organic Fertilizer	10000	175	75	37.5
5	¾ NPK+1 Organic Fertilizer	10000	262.5	112.5	56.25
6	1 NPK+1 Organic Fertilizer	10000	350	150	75
7	¾ NPK + ¼ Organic Fertilizer	2500	262.5	112.5	56.25
8	¾ NPK + ½ Organic Fertilizer	5000	262.5	112.5	56.25
9	¾ NPK + ¾ Organic Fertilizer	7500	262.5	112.5	56.25

Table 2: Characteristics of Inceptisol Soil at the Research Site

S. No.	Parameter	Value	Unit	Rating*
1	N-total	0.28	%	Medium
2	P-available	1.17	ppm	Very Low
3	K-available	0.46	(me/100 g)	Medium
4	C-Organic	1.55	%	Low
5	pH	6.5	-	Somewhat Acidic (5.5-6.5)
6	BV	0.92	(g/cm³)	Medium

Source: Primary data. *Based on the scoring of the Indonesian Soil Research Institute (2023)

Table 3: Analytical results of solid organic fertilizer

S. No.	Parameter	Value
1	Moisture Content (%)	20.08%
2	N-total (%)	1.32%
3	P ₂ O ₅ (%)	2.62%
4	K ₂ O (%)	1.38%
5	pH	7.69
6	C-Organic (%)	22.51%
7	C/N	17.05

Table 4: Average CO₂ emission values at various organic and inorganic fertilizer balances. The highest CO₂ emission values were dominated by the balance treatment with high organic fertilizer doses

Treatment	Soil CO ₂ emission (mg CO ₂ /m ₂ /week)
No fertilizer	1.92 ^c
Standard NPK	4.73 ^b
¼ NPK+1 Organic Fertilizer	5.62 ^a
½ NPK+1 Organic Fertilizer	5.74 ^a
¾ NPK+1 Organic Fertilizer	5.80 ^a
1 NPK+1 Organic Fertilizer	6.16 ^a
¾ NPK + ¼ Organic Fertilizer	4.19 ^b
¾ NPK + ½ Organic Fertilizer	4.72 ^b
¾ NPK + ¾ Organic Fertilizer	5.81 ^a

Different lowercase letters in rows and columns indicate significant differences at 5% DMRT

On the other hand, the application of inorganic fertilizers results in high plant growth (Rahayu *et al.*, 2019; Baharuddin & Sutriana, 2020), including plant roots (Serri *et al.*, 2021; Syamsiyah *et al.*, 2023c), so that root respiration increases and causes CO₂ emissions, this can be seen in the correlation between CO₂ emissions and root biomass ($r=0.330^*$) (Figure 2). These results align with research by Oladele and Adetunji (2021) that applying

Table 5: Correlation between CO₂ emissions and soil properties

	RB	pH	SOC	C-mic
Soil CO ₂ emission	0.330*	0.568**	0.834**	0.527**

RB=Root Biomass, SOC=Soil Organic Carbon, C-mic=C-microba, * and ** indicate significance at $p<0.05$ and $p<0.01$, respectively

Table 6: Average soil C-stock values at different organic and inorganic fertilizer rates. Soil C-stock was lower in the treatment without organic fertilizer addition

Treatment	Soil C-Stock (Mg/ha)
No fertilizer	28.55 ^d
Standard NPK	32.53 ^d
¼ NPK+1 Organic Fertilizer	50.25 ^c
½ NPK+1 Organic Fertilizer	58.61 ^{ab}
¾ NPK+1 Organic Fertilizer	59.13 ^{ab}
1 NPK+1 Organic Fertilizer	64.04 ^a
¾ NPK + ¼ Organic Fertilizer	33.25 ^d
¾ NPK + ½ Organic Fertilizer	50.78 ^c
¾ NPK + ¾ Organic Fertilizer	53.51 ^{bc}

Different lowercase letters in rows and columns indicate significant differences at 5% DMRT




Figure 1: Schematic of alkali trap installation (Modified alkali trap (Singh *et al.*, 2009))




Figure 2: Relationship pattern of CO₂ emission and root biomass in various treatments. Bars and lines represent means ($n=9$) and the same letter appearing on bars and lines throughout the figure indicates no significant difference (according to DMRT test) (Description: Statistical analysis).

Table 7: The average value of plant carbon sequestration in various combinations of organic and inorganic fertilizers. The highest carbon sequestration value was achieved in the balanced treatment with high addition of organic fertilizer

Treatment	Plant Biomass Weight (Kg)			Plant Biomass C Content (%)			Plant Biomass C Sequestration (Mg/ha)*			Plant Carbon Sequestration (Mg/ha)
	Stem	Leaf	Root	Stem	Leaf	Root	Stem	Leaf	Root	
No fertilizer	0.063	0.023	0.083	43.31	4.89	2.83	1.92	0.08	0.16	2.17 ^d
Standard NPK	0.087	0.040	0.110	52.15	6.06	4.90	3.09	0.17	0.38	3.65 ^c
¼ NPK+1 OF	0.120	0.039	0.120	55.01	6.53	5.89	4.78	0.18	0.48	5.44 ^{ab}
½ NPK+1 OF	0.120	0.040	0.130	58.31	6.19	5.76	4.97	0.18	0.55	5.69 ^a
¾ NPK+1 OF	0.123	0.039	0.100	56.16	6.27	5.60	4.94	0.18	0.40	5.51 ^a
1 NPK+1 OF	0.120	0.043	0.100	61.74	6.74	6.13	5.33	0.21	0.45	5.98 ^a
¾ NPK + ¼ OF	0.083	0.033	0.110	52.18	6.19	5.71	3.14	0.15	0.47	3.76 ^c
¾ NPK + ½ OF	0.110	0.037	0.087	54.24	6.50	5.70	4.25	0.17	0.35	4.77 ^b
¾ NPK + ¾ OF	0.123	0.041	0.100	53.53	6.46	6.15	4.66	0.19	0.44	5.29 ^{ab}
Average C Sequestration of Plants (Mg/ha)										4.69

Different lowercase letters in rows and columns indicate significant differences at 5% DMRT. *Biomass C sequestration=Biomass weight x C biomass %

Table 8: The average value of N, P, and K nutrient uptake in various combinations of organic and inorganic fertilizers

Treatment	Nutrient Uptake		
	N (g/plant)	P (g/plant)	K (g/plant)
No fertilizer	1.27 ^f	0.28 ^h	0.86 ^e
Standard NPK	3.17 ^{de}	0.76 ^{fg}	2.03 ^{cd}
¼ NPK+1 Organic Fertilizer	2.91 ^e	0.98 ^{cd}	2.39 ^{ab}
½ NPK+1 Organic Fertilizer	3.43 ^{cd}	1.11 ^{ab}	2.54 ^a
¾ NPK+1 Organic Fertilizer	4.03 ^{ab}	1.07 ^{bc}	2.31 ^{abc}
1 NPK+1 Organic Fertilizer	4.38 ^a	1.22 ^a	2.59 ^a
¾ NPK + ¼ Organic Fertilizer	3.31 ^d	0.68 ^g	1.91 ^d
¾ NPK + ½ Organic Fertilizer	3.73 ^{bc}	0.83 ^{ef}	2.15 ^{bcd}
¾ NPK + ¾ Organic Fertilizer	4.13 ^a	0.94 ^{de}	2.37 ^{abc}

Different lowercase letters in rows and columns indicate significant differences at 5% DMRT

Table 9: Average values of plant growth at various balances of organic and inorganic fertilizers

Treatment	Plant Growth		
	Plant Height (cm)	Number of Leaves (blades)	Stem Diameter (mm)
No fertilizer	151 ^c	12 ^c	2.99 ^c
Standard NPK	180 ^b	14 ^{ab}	3.72 ^{ab}
¼ NPK+1 Organic Fertilizer	183 ^{ab}	13 ^{bc}	3.94 ^{ab}
½ NPK+1 Organic Fertilizer	194 ^{ab}	14 ^{ab}	3.99 ^{ab}
¾ NPK+1 Organic Fertilizer	200 ^a	15 ^{ab}	4.10 ^{ab}
1 NPK+1 Organic Fertilizer	202 ^a	15 ^a	4.16 ^a
¾ NPK + ¼ Organic Fertilizer	198 ^{ab}	13 ^{bc}	3.97 ^{ab}
¾ NPK + ½ Organic Fertilizer	191 ^{ab}	13 ^{bc}	3.90 ^{ab}
¾ NPK + ¾ Organic Fertilizer	191 ^{ab}	13 ^{bc}	3.64 ^b

Different lowercase letters in rows and columns indicate significant differences at 5% DMRT

high N fertilizer can increase plant growth and root biomass, which affects the level of plant root respiration. Root respiration at the ecosystem level is the result of multiplying specific root respiration and root biomass. In other words, the total CO₂ released through root respiration in the ecosystem is directly related to the total amount of root mass in the soil. An increase in root biomass leads to more metabolically active root tissues, thus producing more CO₂ (Jarvi & Burton, 2020). Conversely, a decrease in root biomass will reduce total root respiration due to less root tissue involved in the process. CO₂ emissions were

also influenced by an increase in soil pH ($r=0.568^{**}$) (Table 5), low soil pH can inhibit soil microbial activity. Soil acidity can be reduced by applying soil amendments, where these soil amendments can not only increase soil pH but also play a role in increasing microbial activity which has an impact on increasing heterotrophic respiration and ultimately contributes to increasing soil respiration (Zhang *et al.*, 2019). Soil pH can increase cellulose decomposition by almost twofold due to increased soil microbial activity, leading to more excellent emission release (Moilanen *et al.*, 2012). The combination of ½ NPK and 1 organic fertilizer (10 tons/ha) produced lower CO₂ emissions than 1 NPK and 1 organic fertilizer, although not significantly different, but higher than no fertilizer and standard NPK (Table 4). This is thought to be related to root weight, where this treatment has the highest root weight compared to other treatments.

Efforts to increase carbon storage in soil can reduce CO₂ levels in the atmosphere. The results showed that soil C-Stock content was significantly influenced by organic fertilizer and inorganic fertilizer ($p=0.000$). In the treatment given organic fertilizer input, the soil C-Stock value was higher due to the supply of organic matter (Table 6). This result is consistent with Zhang *et al.* (2022) that the simultaneous application of mineral and organic fertilizers can increase soil C-Organic.

The balance of ½ NPK and 1 organic fertilizer produced soil C-Stock that was not significantly different from 1 NPK and 1 organic fertilizer and higher than no fertilizer and standard NPK (Table 6). These results align with Brar *et al.* (2013) that the application of a combination of organic and inorganic fertilizers with a proportion of 100% NPK + FYM (farmyard manure) significantly increased soil carbon sequestration and was able to increase soil organic carbon content up to 88%. Zuo *et al.* (2023) state that carbon inputs in crop residues and manure are essential in soil organic carbon sequestration, which can directly increase soil C-Stock. Applying organic fertilizers to replace synthetic N fertilizers (with equivalent N levels) can significantly increase annual SOC sequestration by about 700 kg C ha⁻¹ year⁻¹ in upland soils (Xia *et al.*, 2017).

CO₂ emission levels can be reduced by increasing the amount of CO₂ sequestration by plants. In this study, plant carbon

Table 10: The average value of plant carbon balance in various combinations of organic and inorganic fertilizers

Treatment	C- INPUT			C- OUTPUT		Carbon balance (Kg/ha)	
	Plant Biomass C Sequestration (Kg/ha)			Soil C-Stock (Kg/ha)	C-input of fertilizer (Kg/ha)		CO ₂ -C flux (Kg/ha)
	Stem	Leaf	Root				
No fertilizer	1918.06	81.67	163.61	28554.41	0.00	0.22	30717.52 ^d
Standard NPK	3089.67	172.08	386.21	32529.23	0.00	0.54	36176.64 ^d
¼ NPK+1 OF	4781.78	182.05	477.65	50252.25	1125.50	0.64	56818.58 ^c
½ NPK+1 OF	4970.47	175.40	547.65	55648.14	1125.50	0.66	65425.43 ^{ab}
¾ NPK+1 OF	4935.23	175.85	397.37	59131.79	1125.50	0.66	65765.08 ^{ab}
1 NPK+1 OF	5329.69	206.37	446.69	64035.94	1125.50	0.70	71143.49 ^a
¾ NPK + ¼ OF	3138.83	145.25	472.36	33246.12	281.38	0.48	37283.46 ^d
¾ NPK + ½ OF	4245.59	172.31	350.58	50783.75	562.75	0.54	56114.44 ^c
¾ NPK + ¾ OF	4661.83	188.07	437.63	53505.93	844.13	0.66	59636.92 ^{bc}

Different lowercase letters in rows and columns indicate significant differences at 5% DMRT

sequestration was significantly affected by organic fertilizer and inorganic fertilizer ($p=0.000$). This can be caused by an increase in soil fertility conditions due to an increase in organic carbon levels in the soil, where organic carbon levels act as a source of nutrients (Hairiah & Rahayu, 2007; Masryfah *et al.*, 2019; Punuindoong *et al.*, 2021) which affects plant growth (Purakayastha *et al.*, 2008). Plant carbon sequestration in various treatments ranged from 2.10-5.98 Mg/ha (Table 7). The experimental results showed that the treatment of ½ NPK and 1 organic fertilizer resulted in carbon sequestration, which was not significantly different from 1 NPK and 1 organic fertilizer and higher than no fertilizer and standard NPK. These results align with Sulaeman *et al.* (2016) that the input of organic fertilizer plus NPK fertilizer at a dose of 50 and 75% of the recommended dose can produce an increase in plant growth manifested in total biomass production and significantly increase carbon sequestration in biomass compared to the use of organic fertilizer alone. This can be seen in the correlation between total plant biomass and plant height ($r=0.612^{**}$), number of leaves ($r=0.423^{*}$), and stem diameter ($r=0.548^{**}$).

The balance also achieved higher plant uptake of N, P, and K nutrients compared to no fertilizer and standard NPK (Table 8). The reduction of inorganic fertilizers balanced with organic fertilizers can increase the value of N-total, P-available, K-available in the soil so as to increase the uptake of N, P, and K nutrients in plant tissues (Table 9) (Syamsiyah *et al.*, 2023d), followed by increased plant growth where the element that plays the most role in increasing plant growth, especially plant height, is nitrogen nutrients ($r=0.816^{**}$) (Syamsiyah *et al.*, 2020). Maize plants need nitrogen nutrients both in the vegetative growth phase and the generative phase. Nitrogen is a constituent in the hormone auxin, a plant growth hormone (Oleńska *et al.*, 2020). Plant physiological processes will produce height and width in plants, affecting the number of leaves ($r=0.582^{**}$) (Marian & Tuhuteru, 2019). In addition, the results showed a positive correlation between stem diameter and potassium levels in plant tissue ($r=0.721^{**}$). Potassium is an essential element that plays a vital role in increasing stem diameter due to its role in increasing the level of sclerenchyma in the stem, which provides thickening and strength to the stem tissue (Attia *et al.*, 2022). This also explains why the balanced treatment of organic and inorganic fertilizers in this study had higher plant C sequestration results than the treatment without fertilizer and standard NPK.

Carbon balance is calculated using all carbon inputs and outputs (Rutledge *et al.*, 2015). The soil's supply or carbon input can be achieved from photosynthesis and other organic matter from outside. Carbon output can be affected by the decomposition process due to the conversion of soil carbon into CO₂ (Liddicoat *et al.*, 2010). In this study, organic and inorganic fertilizers significantly affected the carbon balance ($p=0.000$). The average carbon balance of all treatments ranged from 30717.52-71143.49 Kg/ha (Table 10). The combination of ½ NPK and 1 organic fertilizer (10 tons/ha) produced a lower carbon balance than 1 NPK and 1 organic fertilizer, although not significantly different, but higher than no fertilizer and standard NPK (Table 10). The carbon balance results show that applying organic fertilizer causes higher CO₂ storage than CO₂ released. Fertile soil conditions accompanied by high plant productivity allow carbon input through photosynthesis to exceed the amount of carbon lost (Siringoringo, 2014; Persiani *et al.*, 2019). These results align with Xu *et al.* (2020) that tillage practices that can increase carbon inputs, such as applying organic fertilizers, are highly recommended because of their high efficiency in supporting soil carbon sequestration processes.

CONCLUSIONS AND SUGGESTION

The use of a fertilizer combination of NPK and organic fertilizers significantly increased soil carbon stocks (33.25-64.04 Mg/ha) and carbon sequestration (3.76-5.98 Mg/ha). The integration of organic fertilizers with inorganic fertilizers has proven to be effective in increasing C sequestration, reducing production costs, and increasing agricultural productivity. Thus, the use of organic fertilizers in agricultural systems is essential to improve fertilizer use efficiency, maintain carbon balance, and reduce negative impacts on the environment.

ACKNOWLEDGMENT

The authors would like to thank God for making all this possible. The authors would also like to thank the Department of Soil Science, Sebelas Maret University, for supporting funds in conducting the research and the laboratory assistants of the Soil Chemistry and Fertility Laboratory of Sebelas Maret University for their kind cooperation in the data collection process.

REFERENCES

- Abera, W., Tamene, L., Abegaz, A., Hailu, H., Piikki, K., Söderström, M., Girvetz, E., & Sommer, R. (2021). Estimating spatially distributed SOC sequestration potentials of sustainable land management practices in Ethiopia. *Journal of Environmental Management*, 286, 112191. <https://doi.org/10.1016/j.jenvman.2021.112191>
- Al-Ghussain, L. (2019). Global warming: review on driving forces and mitigation. *Environmental Progress and Sustainable Energy*, 38(1), 13-21. <https://doi.org/10.1002/ep.13041>
- Attia, H., Rebah, F., Ouhibi, C., Saleh, M. A., Althobaiti, A. T., Alamer, K. H., Nasri, M. B., & Lachaâl, M. (2022). Effect of Potassium Deficiency on Physiological Responses and Anatomical Structure of Basil, *Ocimum basilicum* L. *Biology*, 11(11), 1-14. <https://doi.org/10.3390/biology11111557>
- Baharuddin, R., & Sutriana, S. (2020). Growth and Production of Tumpangsari Chili Plants with Red Onion Through and NPK Fertilization in Peat Raisa. *Dinamika Pertanian*, 35(3), 73-80. [https://doi.org/10.25299/dp.2019.vol35\(3\).4567](https://doi.org/10.25299/dp.2019.vol35(3).4567)
- Béziat, P., Ceschia, E., & Dedieu, G. (2009). Carbon balance of a three crop succession over two cropland sites in South West France. *Agricultural and Forest Meteorology*, 149(10), 1628-1645. <https://doi.org/10.1016/j.agrformet.2009.05.004>
- Brar, B. S., Singh, K., Dheri, G. S., & Balwinder-Kumar. (2013). Carbon sequestration and soil carbon pools in a rice-wheat cropping system: Effect of long-term use of inorganic fertilizers and organic manure. *Soil and Tillage Research*, 128, 30-36. <https://doi.org/10.1016/j.still.2012.10.001>
- Chauhan, B. S., Prabhjyot-Kaur, Mahajan, G., Randhawa, R. K., Singh, H., & Kang, M. S. (2014). Global Warming and Its Possible Impact on Agriculture in India. *Advances in Agronomy*, 123, 65-121. <https://doi.org/10.1016/B978-0-12-420225-2.00002-9>
- Chu, H., Hosen, Y., & Yagi, K. (2007). NO, N₂O, CH₄ and CO₂ fluxes in winter barley field of Japanese Andisol as affected by N fertilizer management. *Soil Biology and Biochemistry*, 39(1), 330-339. <https://doi.org/10.1016/j.soilbio.2006.08.003>
- de Urzedo, D. I., Franco, M. P., Pitombo, L. M., & do Carmo, J. B. (2013). Effects of organic and inorganic fertilizers on greenhouse gas (GHG) emissions in tropical forestry. *Forest Ecology and Management*, 310, 37-44. <https://doi.org/10.1016/j.foreco.2013.08.018>
- Dossou-Yovo, E. R., Brüggemann, N., Ampofo, E., Igwe, A. M., Jesse, N., Huat, J., & Agbossou, E. K. (2016). Combining no-tillage, rice straw mulch and nitrogen fertilizer application to increase the soil carbon balance of upland rice field in northern Benin. *Soil and Tillage Research*, 163, 152-159. <https://doi.org/10.1016/j.still.2016.05.019>
- Ebrahimi, M., Sarikhani, M. R., Sinegani, A. A. S., Ahmadi, A., & Keesstra, S. (2019). Estimating the soil respiration under different land uses using artificial neural network and linear regression models. *Catena*, 174, 371-382. <https://doi.org/10.1016/j.catena.2018.11.035>
- Ekowati, D., & Nasir, M. (2011). The Growth of Maize Crop (*Zea mays* L.) BSI-2 Variety on Rejected and non Rejected Sand at Pantai Trisik Kulon Progo. *Jurnal Manusia Dan Lingkungan*, 18(3), 220-231.
- Ferrini, F., Fini, A., Mori, J., & Gori, A. (2020). Role of vegetation as a mitigating factor in the urban context. *Sustainability*, 12(10), 4247. <https://doi.org/10.3390/su12104247>
- Franzluebbers, A. J., & Veum, K. S. (2020). Comparison of two alkali trap methods for measuring the flush of CO₂. *Agronomy Journal*, 112(2), 1279-1286. <https://doi.org/10.1002/agj2.20141>
- Hairiah, K., & Rahayu, S. (2007). *Measurement of Carbon Stored in Different Land Uses*. Nairobi, Kenya: World Agroforestry Centre.
- IPCC. (2006). *IPCC Guidelines for National Greenhouse Gas Inventories*. Prepared by the National Greenhouse Gas Inventories Programme, H. S. Eggleston, K. Miwa, N. Srivastava & K. Tanabe (Eds.). Japan: IGES.
- Irawan, Maswar, Yustika, R. D., & Ariani, R. (2002). *Sifat Fisika Tanah dan Metode Analisisnya*. Java, Indonesia: Balai Penelitian Tanah.
- Jarvi, M. P., & Burton, A. J. (2020). Root respiration and biomass responses to experimental soil warming vary with root diameter and soil depth. *Plant and Soil*, 451, 435-446. <https://doi.org/10.1007/s11104-020-04540-1>
- Kartikawati, R., & Nursyamsi, D. (2013). Pengaruh Pengairan, Pemupukan, Dan Penghambat Nitrifikasi Terhadap Emisi Gas Rumah Kaca Di Lahan Sawah Tanah Mineral. *Ecolab*, 7(2), 93-107.
- Li, H., Feng, W., He, X., Zhu, P., Gao, H., Sin, N., & Xu, M. (2017). Chemical fertilizers could be completely replaced by manure to maintain high maize yield and soil organic carbon (SOC) when SOC reaches a threshold in the Northeast China Plain. *Journal of Integrative Agriculture*, 16(4), 937-946. [https://doi.org/10.1016/S2095-3119\(16\)61559-9](https://doi.org/10.1016/S2095-3119(16)61559-9)
- Li, L.-J., You, M.-Y., Shi, H.-A., Ding, X.-L., Qiao, Y.-F., & Han, X.-Z. (2013). Soil CO₂ emissions from a cultivated Mollisol: Effects of organic amendments, soil temperature, and moisture. *European Journal of Soil Biology*, 55, 83-90. <https://doi.org/10.1016/j.ejsobi.2012.12.009>
- Liang, Z., Jin, X., Zhai, P., Zhao, Y., Cai, J., Li, S., Yang, S., Li, C., & Li, C. (2022). Combination of organic fertilizer and slow-release fertilizer increases pineapple yields, agronomic efficiency and reduces greenhouse gas emissions under reduced fertilization conditions in tropical areas. *Journal of Cleaner Production*, 343, 131054. <https://doi.org/10.1016/j.jclepro.2022.131054>
- Liddicoat, C., Schapel, A., Davenport, D., & Dwyer, E. (2010). *Soil carbon and climate change*. PIRSA Discussion Paper.
- Liu, D., Liu, X., Liu, Y., Li, L., Pan, G., Crowley, D., & Tippkötter, R. (2011). Soil organic carbon (SOC) accumulation in rice paddies under long-term agro-ecosystem experiments in southern China – VI. Changes in microbial community structure and respiratory activity. *Biogeosciences Discussions*, 8, 1529-1554. <https://doi.org/10.5194/bgd-8-1529-2011>
- Liu, H., Zhou, G., Bai, S. H., Song, J., Shang, Y., He, M., Wang, X., & Zheng, Z. (2019). Differential response of soil respiration to nitrogen and phosphorus addition in a highly phosphorus-limited subtropical forest, China. *Forest Ecology and Management*, 448, 499-508. <https://doi.org/10.1016/j.foreco.2019.06.020>
- Machado, C. B., Lima, J. R. D. S., Antonino, A. C. D., de Souza, E. S., Souza, R. M. S., & Alves, E. M. (2016). Daily and seasonal patterns of CO₂ fluxes and evapotranspiration in maize-grass intercropping. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 20(9), 777-782. <https://doi.org/10.1590/1807-1929/agriambi.v20n9p777-782>
- Marian, E., & Tuhuteru, S. (2019). Utilization of Tofu Liquid Waste to Growth and Yield of Chicory (*Brassica pekinensis*). *Agritrop*, 17(2), 134-144. <https://doi.org/10.32528/agritrop.v17i2.2663>
- Masryfah, T. F., Banuwa, I. S., Buchari, H., & Utomo, M. (2019). Effects of tillage system and herbicide application on aboveground carbon stored in maize (*Zea mays* L.) 3rd cropping season. *Jurnal Agrotek Tropika*, 7(3), 433-441. <https://doi.org/10.23960/jat.v7i3.3547>
- Maswar, M., Husnain, H., & Agus, F. (2014). Effect of Manure and Chemical Fertilizers on CO₂ Fluxes on Drained Peatlands. *Jurnal Tanah Dan Iklim*, 38(2), 71-77.
- Moilanen, M., Hytönen, J., & Leppälä, M. (2012). Application of wood ash accelerates soil respiration and tree growth on drained peatland. *European Journal of Soil Science*, 63(4), 467-475. <https://doi.org/10.1111/j.1365-2389.2012.01467.x>
- Nema, P., Nema, S., & Roy, P. (2012). An overview of global climate changing in current scenario and mitigation action. *Renewable and Sustainable Energy Reviews*, 16(4), 2329-2336. <https://doi.org/10.1016/j.rser.2012.01.044>
- Oladele, S. O., & Adetunji, A. T. (2021). Agro-residue biochar and N fertilizer addition mitigates CO₂-C emission and stabilized soil organic carbon pools in a rain-fed agricultural cropland. *International Soil and Water Conservation Research*, 9(1), 76-86. <https://doi.org/10.1016/j.iswcr.2020.09.002>
- Oleńska, E., Małek, W., Wójcik, M., Swiecicka, I., Thijs, S., & Vangronsveld, J. (2020). Beneficial features of plant growth-promoting rhizobacteria for improving plant growth and health in challenging conditions: A methodical review. *Science of the Total Environment*, 743, 140682. <https://doi.org/10.1016/j.scitotenv.2020.140682>
- Ölinger, R., Beck, T., Heilmann, B., & Beese, F. (1996). Soil Respiration. In F. Schinner, R. Öhlinger, E. Kandeler & R. Margesin (Eds.), *Methods in Soil Biology* (pp. 93-110) Berlin, Heidelberg: Springer. https://doi.org/10.1007/978-3-642-60966-4_6
- Persiani, A., Diacono, M., Monteforte, A., & Montemurro, F. (2019). Agronomic performance, energy analysis, and carbon balance comparing different fertilization strategies in horticulture under Mediterranean conditions. *Environmental Science and Pollution Research*, 26, 19250-19260. <https://doi.org/10.1007/s11356-019-05292-x>
- Punuindoong, S., Sinolungan, M. T. M., & Rondonuwu, J. J. (2021). Kajian Nitrogen, Fosfor, Kalium dan C-Organik Pada Tanah Berpasir Pertanian Kelapa Desa Ranoketang Atas. *Soil and Environment Journal*, 1(1), 6-11.

- Purakayastha, T. J., Rudrappa, L., Singh, D., Swarup, A., & Bhadraray, S. (2008). Long-term impact of fertilizers on soil organic carbon pools and sequestration rates in maize-wheat-cowpea cropping system. *Geoderma*, 144(1-2), 370-378. <https://doi.org/10.1016/j.geoderma.2007.12.006>
- Rahayu, Syamsiyah, J., Cahyani, V. R., & Fauziah, S. K. (2019). The effects of biochar and compost on different cultivars of shallots (*Allium ascalonicum* L.) growth and nutrient uptake in sandy soil under saline water. *Journal of Soil Science and Agroclimatology*, 16(2), 216-228. <https://doi.org/10.20961/STJSSA.V16I2.34209>
- Rasse, D. P., Rumpel, C., & Dignac, M.-F. (2005). Is soil carbon mostly root carbon? Mechanisms for a specific stabilisation. *Plant and Soil*, 269, 341-356. <https://doi.org/10.1007/s11104-004-0907-y>
- Romadhan, P., Gusmini, & Hermansah. (2022). Improvement of The Chemical Properties of the Ex-Gold Mining Land Through the Application of Organic Fertilizer Granular Biokanat. *Journal of Agricultural Science*, 20(1), 74-83.
- Roß, C.-L., Baumecker, M., Ellmer, F., & Kautz, T. (2022). Organic Manure Increases Carbon Sequestration Far beyond the "4 per 1000 Initiative" Goal on a Sandy Soil in the Thyrow Long-Term Field Experiment DIV2. *Agriculture*, 12(2), 170. <https://doi.org/10.3390/agriculture12020170>
- Rotbart, N., Borisover, M., Bukhanovsky, N., Nasonova, A., Bar-Tal, A., & Oren, A. (2017). Examination of residual chloroform interference in the measurement of microbial biomass C by fumigation-extraction. *Soil Biology and Biochemistry*, 111, 60-65. <https://doi.org/10.1016/j.soilbio.2017.03.018>
- Russell, A. E., Cambardella, C. A., Laird, D. A., Jaynes, D. B., & Meek, D. W. (2009). Nitrogen fertilizer effects on soil carbon balances in Midwestern U.S. agricultural systems. *Ecological Applications*, 19(5), 1102-1113. <https://doi.org/10.1890/07-1919.1>
- Rutledge, S., Mudge, P. L., Campbell, D. I., Woodward, S. L., Goodrich, J. P., Wall, A. M., Kirschbaum, M. U. F., & Schipper, L. A. (2015). Carbon balance of an intensively grazed temperate dairy pasture over four years. *Agriculture, Ecosystems & Environment*, 206, 10-20. <https://doi.org/10.1016/j.agee.2015.03.011>
- Sainju, U. M., Ghimire, R., & Dangi, S. (2021). Soil carbon dioxide and methane emissions and carbon balance with crop rotation and nitrogen fertilization. *Science of the Total Environment*, 775, 145902. <https://doi.org/10.1016/j.scitotenv.2021.145902>
- Serri, F., Souri, M. K., & Rezapanah, M. (2021). Growth, biochemical quality and antioxidant capacity of coriander leaves under organic and inorganic fertilization programs. *Chemical and Biological Technologies in Agriculture*, 8, 33. <https://doi.org/10.1186/s40538-021-00232-9>
- Shakoor, A., Dar, A. A., Arif, M. S., Farooq, T. H., Yasmeen, T., Shahzad, S. M., Tufail, M. A., Ahmed, W., Albasher, G., & Ashraf, M. (2022). Do soil conservation practices exceed their relevance as a countermeasure to greenhouse gases emissions and increase crop productivity in agriculture? *Science of the Total Environment*, 805, 150337. <https://doi.org/10.1016/j.scitotenv.2021.150337>
- Singh, K. P., Ghoshal, N., & Singh, S. (2009). Soil carbon dioxide flux, carbon sequestration and crop productivity in a tropical dryland agroecosystem: Influence of organic inputs of varying resource quality. *Applied Soil Ecology*, 42(3), 243-253. <https://doi.org/10.1016/j.apsoil.2009.04.005>
- Siringoringo, H. H. (2014). The important Roles of managing carbon sequestration in soils. *Jurnal Analisis Kebijakan Kehutanan*, 11(2), 175-192.
- Sulaeman, Y., Maswar, & Erfandi, D. (2016). Effect of Organic and Inorganic Fertilizers Combination on Soil Productivity and Crop Yield of Maize Farming in Acid Upland Soil. *Jurnal Pengkajian Dan Pengembangan Teknologi Pertanian*, 20(1), 1-12.
- Suwandi, Sopha, G., & Yufdy, M. (2015). The Effectiveness of Organic Fertilizer, NPK, and Biofertilizer Managements on Growth and Yields of Shallots. *Jurnal Hortikultura*, 25(3), 208-221.
- Syamsiyah, J., Ariyanto, D. P., Herawati, A., Herdiansyah, G., & Dwisetio, P. K. (2023a). Rice paddy field on Gentungan, Karanganyar: Soil C Humic Acid, Fulvic Acid, and Stock as Affected by Period of Organic Practices. *IOP Conference Series: Earth and Environmental Science*, 1165, 012013. <https://doi.org/10.1088/1755-1315/1165/1/012013>
- Syamsiyah, J., Ariyanto, D. P., Komariah, Herawati, A., Dwisetio, P. K., Sari, S. I., Salsabila, H. A., Herdiansyah, G., Hartati, S., & Mujiyo. (2023b). Temporal variation in the soil properties and rice yield of organic rice farming in the tropical monsoon region, Indonesia. *Journal of Soil Science and Agroclimatology*, 20(2), 231-239. <https://doi.org/10.20961/stjssa.v20i2.71431>
- Syamsiyah, J., Minardi, S., Herdiansyah, G., Cahyono, O., & Mentari, F. C. (2023c). Physical Properties of Alfisols, Growth and Products of Hybrid Corn Affected by Organic and Inorganic Fertilizer. *Journal of Sustainable Agriculture*, 38(1), 99-112. <https://doi.org/10.20961/carakatani.v38i1.65014>
- Syamsiyah, J., Minardi, S., Khadaffi, J., Hartati, S., & Herdiansyah, G. (2023d). Partial substitution inorganic using organic fertilizers on soil N, P, K and corn yield on inceptisols. *Jurnal AGRO*, 10(2), 242-251. <https://doi.org/10.15575/27875>
- Syamsiyah, J., Rahayu, R., & Binafsihi, W. (2020). Soil properties and shallot yield responses to different salinity levels. *Journal of Soil Science and Agroclimatology*, 17(1), 30-34. <https://doi.org/10.20961/stjssa.v17i1.41566>
- Syamsiyah, J., Sunarminto, B. H., Hanudin, E., Widada, J., & Mujiyo. (2019). Carbon dioxide emission and carbon sequestration potential in Alfisol. *Bulgarian Journal of Agricultural Science*, 25(1), 42-48.
- Wang, J., Xie, J., Li, L., Effah, Z., Xie, L., Luo, Z., Zhou, Y., & Jiang, Y. (2022). Fertilization treatments affect soil CO₂ emission through regulating soil bacterial community composition in the semiarid Loess Plateau. *Scientific Reports*, 12(1), 1-11. <https://doi.org/10.1038/s41598-022-21108-4>
- Wang, Y., Hu, C., Dong, W., Li, X., Zhang, Y., Qin, S., & Oenema, O. (2015). Carbon budget of a winter-wheat and summer-maize rotation cropland in the North China plain. *Agriculture, Ecosystems and Environment*, 206, 33-45. <https://doi.org/10.1016/j.agee.2015.03.016>
- Wu, W., Ma, X., Zhang, Y., Li, W., & Wang, Y. (2020). A novel conformable fractional non-homogeneous grey model for forecasting carbon dioxide emissions of BRICS countries. *Science of the Total Environment*, 707, 135447. <https://doi.org/10.1016/j.scitotenv.2019.135447>
- Xia, L., Lam, S. K., Yan, X., & Chen, D. (2017). How Does Recycling of Livestock Manure in Agroecosystems Affect Crop Productivity, Reactive Nitrogen Losses, and Soil Carbon Balance? *Environmental Science and Technology*, 51(13), 7450-7457. <https://doi.org/10.1021/acs.est.6b06470>
- Xu, M., & Qi, Y. (2001). Soil-surface CO₂ efflux and its spatial and temporal variations in a young ponderosa pine plantation in northern California. *Global Change Biology*, 7(6), 667-677. <https://doi.org/10.1046/j.1354-1013.2001.00435.x>
- Xu, S., Sheng, C., & Tian, C. (2020). Changing soil carbon: Influencing factors, sequestration strategy and research direction. *Carbon Balance and Management*, 15(1), 1-9. <https://doi.org/10.1186/s13021-020-0137-5>
- Yuan, J., Sha, Z., Hassani, D., Zhao, Z., & Cao, L. (2017). Assessing environmental impacts of organic and inorganic fertilizer on daily and seasonal Greenhouse Gases effluxes in rice field. *Atmospheric Environment*, 155, 119-128. <https://doi.org/10.1016/j.atmosenv.2017.02.007>
- Zhang, C., Zhao, Z., Li, F., & Zhang, J. (2022). Effects of Organic and Inorganic Fertilization on Soil Organic Carbon and Enzymatic Activities. *Agronomy*, 12(12), 3125. <https://doi.org/10.3390/agronomy12123125>
- Zhang, J., Li, Q., Wu, J., & Song, X. (2019). Effects of nitrogen deposition and biochar amendment on soil respiration in a *Torreya grandis* orchard. *Geoderma*, 355, 113918. <https://doi.org/10.1016/j.geoderma.2019.113918>
- Zhang, L., Zhuang, Q., He, Y., Liu, Y., Yu, D., Zhao, Q., Shi, X., Xing, S., & Wang, G. (2016). Toward optimal soil organic carbon sequestration with effects of agricultural management practices and climate change in Tai-Lake paddy soils of China. *Geoderma*, 275, 28-39. <https://doi.org/10.1016/j.geoderma.2016.04.001>
- Zulkarnain, M., Prasetya, B., & Soemarno. (2013). Pengaruh kompos, pupuk kandang, dan custom-bio terhadap sifat tanah, pertumbuhan dan hasil tebu (*Saccharum officinarum* L.) pada entisol di kebun ngrangkah-pawon, kediri). *Indonesian Green Technology Journal*, 2(1), 45-52.
- Zuo, W., Gu, B., Zou, X., Peng, K., Shan, Y., Yi, S., Shan, Y., Gu, C., & Bai, Y. (2023). Soil organic carbon sequestration in croplands can make remarkable contributions to China ' s carbon neutrality. *Journal of Cleaner Production*, 382, 135268. <https://doi.org/10.1016/j.jclepro.2022.135268>