



Diversity analysis of sesame germplasm using DIVA-GIS

B Spandana, N Sivaraj¹, G John Prasanna Rao², G Anuradha, S Sivaramakrishnan & Farzana Jabeen³

Institute of Biotechnology (IBT),
Acharya N.G. Ranga Agricultural University, Hyderabad-500 030, India.
E-mail: spandanabandila@gmail.com

Received 07 December 2011; Revised 10 April 2012; Accepted 26 June 2012

Abstract

Sesame (*Sesamum indicum* L.) was studied for its distribution and diversity in India using DIVA-GIS. Grid maps were generated for diversity analysis of the eight quantitative traits viz., plant height, inter-node length, leaves per plant, number of flowers per plant, number of capsules per plant, number of seeds per capsule, seed weight and seed yield. The results indicated that diverse accessions for all these traits can be sourced from Maharashtra, Gujarat and Madhya Pradesh (partly covering Chattisgarh) states and these states are diversity rich pockets for sesame germplasm in India.

Keywords: DIVA-GIS, germplasm, India, mapping, *Sesamum indicum*

Introduction

Sesame (*Sesamum indicum* L.) family Pedaliaceae, is an important source of edible oil and is widely used as one of the ingredients in food products especially in bakery foods and animal feed. Sesame germplasm has tremendous variability. Efficient and systematic exploitation of this diversity is the key to any crop improvement program. This, however, requires as a first step, the assessment of genetic diversity and population structure of the species under consideration. DIVA-GIS, a Geographic Information System (GIS) is designed to map the range of distribution of species in which one is interested (Hijmans *et al.* 2000). GIS has

been successfully used in identifying areas of rich diversity in various parts of the world. DIVA-GIS is a software tool for the analysis of diversity and it enables us to understand and comprehend the distribution of diversity on the geographical scale and also helps in identifying gaps in collection.

Materials and methods

The experiment was carried out during late Kharif 2008–09. The experimental materials were sown in simple Randomised Block Design with 60 × 10 cm spacing in three replications at College Farm, College of Agriculture, Acharya N.G. Ranga Agricultural University, Rajendranagar, Hyderabad. Recommended

¹National Bureau of Plant Genetic Resources (NBPGR), Hyderabad, India.

²Directorate of Rice Research (DRR), Hyderabad-500 030, India.

³Department of Genetics and Plant breeding (ANGRAU), Hyderabad-500 030, India.

agronomic practices and prophylactic measures were adopted.

Assessment of diversity in sesame germplasm and geographical distribution using DIVA-GIS

Characterization data recorded on the sixty accessions of sesame such as plant height, internode length, leaves plant⁻¹, number of flowers plant⁻¹, number of capsules plant⁻¹, number of seeds capsule⁻¹, seed yield and seed weight was subjected to DIVA-GIS analysis. To know the spatial distribution and assessment of variability DIVA-GIS version 7.1.6, was used. Geographical coordinates of the collection sites from various parts of India were also obtained using Garmin 12 Global Positioning System (GPS). India shape file was used for plotting the georeferenced points using the layer menu on the software. Point- to- grid option using 'simple' method on the "Analysis Menu" and the output variables "Diversity and Statistics" were selected for getting the output files. Under diversity, Shannon diversity index was picked and for the statistics, coefficient of variation was selected. Grid maps on the diversity, and coefficient of variation were generated for various traits recorded on the sesame germplasm.

Shannon Diversity Index

DIVA-GIS can calculate a number of different diversity indices for each grid cell. Shannon diversity index was selected for diversity index analysis. The formula for the index taken is given below.

$$\text{Shannon } H' = -\sum pi \ln pi$$

ni – number of individuals in the i -th class

pi – proportional abundance of the i th class = ni / N

Results and discussion

Sesame germplasm (60 accessions) was collected by National Bureau of Plant Genetic Resources, New Delhi was used for studying the variability. The germplasm accessions originated from eight Indian states covering Gujarat (8), Himachal Pradesh (1), Madhya Pradesh (24), Nagaland (1), Punjab (92), Rajasthan (8) and Uttar Pradesh (3). The geographical coordinates of

the sesame collection sites, accession identity and the plant traits used for DIVA-GIS analysis are provided in Table 1. The collection sites were mapped using DIVA-GIS (Fig.1).




Fig 1. DIVA-GIS map showing geographical origin of sesame germplasm used in the study

Observations on eight traits *viz.*, plant height, internode length, leaves plant⁻¹, number of flowers plant⁻¹, number of capsules plant⁻¹, number of seeds capsule⁻¹, seed yield and seed weight were recorded. All quantitative characters exhibited variability evident by high CV observed in most of the traits studied. The high CV observed among morphological characters is an indication of the level of diversity within the populations of sesame found growing in these habitats. Moderately low CV values were observed for internode length and for test seed weight.

Diversity Analysis (Shannon diversity index)

Grid maps were generated for diversity analysis of the eight quantitative traits (Fig. 2, 3 & 4) indicated that diverse accessions for all these traits can be sourced from Maharashtra, Gujarat and Madhya Pradesh (partly covering Chattisgarh) states of India. However, highly diverse sesame germplasm accessions for leaves plant⁻¹ was sourced from Maharashtra and

Table 1. Observations on the plant traits used as input in the DIVA-GIS to assess the geographical distribution of diversity in Sesame germplasm

| Acc Id | State | Latitude | Longitude | capsules/ plant | Plant height | Internode length | leaves/ Plant | Flowers/ plant | Seed weight | Seed yield | Seeds/ capsule |
|---------|------------------|----------|-----------|-----------------|--------------|------------------|---------------|----------------|-------------|------------|----------------|
| IC14163 | Gujarat | 21.6677 | 73.5195 | 55.4 | 25.2 | 3.2 | 103.5 | 50.0 | 3.02 | 6.9 | 40.8 |
| IC43169 | Gujarat | 22.804 | 73.6183 | 23.2 | 23.8 | 4.2 | 43.2 | 25.3 | 2.6 | 3.612 | 59.4 |
| IC43171 | Gujarat | 22.9028 | 72.3831 | 28.0 | 21.0 | 3.6 | 39.6 | 35.6 | 2.9 | 3.66 | 37.3 |
| IC43177 | Gujarat | 23.0017 | 74.0135 | 70.6 | 26.4 | 4.2 | 105.6 | 71.0 | 2.3 | 4.5 | 28.2 |
| IC43179 | Gujarat | 23.5945 | 70.8516 | 47.8 | 24.6 | 3.5 | 68.2 | 38.6 | 2.92 | 4.77 | 52.4 |
| IC43181 | Gujarat | 22.6558 | 71.5926 | 56.8 | 23.5 | 4.0 | 125.4 | 57.0 | 2.9 | 8.45 | 30.4 |
| IC43185 | Gujarat | 24.1874 | 70.4069 | 43.2 | 26.6 | 3.6 | 67.8 | 41.6 | 2.4 | 5.36 | 77.0 |
| IC43217 | Gujarat | 23.7427 | 72.2349 | 38.2 | 23.6 | 3.2 | 55.2 | 39.2 | 2.6 | 5.2 | 53.0 |
| IC96079 | Himachal Pradesh | 31.7959 | 76.7309 | 29.0 | 14.3 | 2.6 | 60.3 | 29.3 | 2.6 | 4.97 | 66.0 |
| IC14329 | Madhya Pradesh | 20.482 | 81.0786 | 33.6 | 23.2 | 3.8 | 82.0 | 35.4 | 2.6 | 5.55 | 63.6 |
| IC21705 | Madhya Pradesh | 21.6183 | 80.8809 | 36.0 | 26.8 | 4.0 | 60.0 | 37.2 | 2.8 | 4.14 | 40.6 |
| IC23233 | Madhya Pradesh | 22.0135 | 83.2524 | 56.0 | 26.6 | 4.0 | 55.0 | 57.2 | 2.8 | 8.4 | 54.0 |
| IC23271 | Madhya Pradesh | 23.2487 | 82.6101 | 35.2 | 30.4 | 4.0 | 67.2 | 35.8 | 2.9 | 7.59 | 74.4 |
| IC23321 | Madhya Pradesh | 20.804 | 75.0511 | 41.8 | 31.6 | 4.0 | 74.6 | 42.6 | 2.8 | 5.3 | 45.8 |
| IC23325 | Madhya Pradesh | 25.719 | 77.8178 | 46.0 | 27.3 | 4.0 | 68.8 | 46.8 | 2.9 | 4.7 | 35.8 |
| IC23327 | Madhya Pradesh | 24.0886 | 77.966 | 52.3 | 30.6 | 4.0 | 115 | 53.6 | 2.7 | 8.33 | 33.5 |
| IC23332 | Madhya Pradesh | 23.5945 | 80.3869 | 19.5 | 25.5 | 3.5 | 48.5 | 20.0 | 2.85 | 2.77 | 50.0 |
| IC23335 | Madhya Pradesh | 18.9998 | 81.4738 | 33.2 | 24.5 | 4.0 | 77.6 | 34.2 | 2.17 | 3.84 | 53.4 |
| IC23341 | Madhya Pradesh | 23.0017 | 78.8059 | 40.5 | 24.2 | 3.9 | 65.7 | 41.4 | 2.79 | 5.845 | 48.0 |
| IC23346 | Madhya Pradesh | 24.1874 | 80.041 | 51.4 | 17.0 | 4.0 | 51.3 | 52.6 | 2.06 | 2.75 | 26.0 |
| IC41932 | Madhya Pradesh | 22.9522 | 77.1261 | 39.4 | 27.6 | 4.0 | 100.8 | 40.2 | 2.7 | 5.06 | 47.6 |
| IC41948 | Madhya Pradesh | 22.31 | 78.7071 | 17.2 | 15.8 | 3.8 | 50.8 | 17.8 | 2.6 | 2.1 | 47.0 |
| IC41953 | Madhya Pradesh | 21.8159 | 75.2981 | 30.4 | 20.5 | 3.8 | 72.6 | 31.2 | 2.7 | 2.7 | 33.0 |
| IC41962 | Madhya Pradesh | 22.557 | 76.4344 | 55.2 | 25.2 | 4.0 | 111.2 | 56.2 | 2.74 | 4.82 | 31.9 |
| IC41964 | Madhya Pradesh | 20.4325 | 81.1774 | 39.0 | 22.5 | 4.0 | 125.0 | 40.0 | 2.9 | 2.34 | 55.0 |
| IC41966 | Madhya Pradesh | 18.8022 | 82.0173 | 31.0 | 21.6 | 4.0 | 53.2 | 31.6 | 2.62 | 3.7 | 43.4 |
| IC42200 | Madhya Pradesh | 20.482 | 82.4125 | 51.4 | 23.3 | 3.6 | 87.8 | 52.2 | 2.6 | 8.04 | 60.2 |
| IC52585 | Madhya Pradesh | 22.804 | 81.3256 | 47.2 | 25.8 | 3.5 | 81.4 | 48.8 | 2.6 | 5.3 | 43.2 |

Cont....

| | | | | | | | | | | | |
|---------|----------------|---------|---------|------|------|-----|-------|------|------|-------|------|
| IC52586 | Madhya Pradesh | 23.9898 | 81.622 | 42.0 | 15.3 | 4.2 | 68.4 | 44.6 | 2.6 | 6.77 | 62.0 |
| IC52592 | Madhya Pradesh | 23.2487 | 78.3612 | 21.5 | 15.6 | 4.2 | 45.5 | 21.6 | 2.7 | 3.5 | 62.0 |
| IC52593 | Madhya Pradesh | 23.7427 | 83.45 | 23.2 | 21.1 | 4.0 | 57.8 | 23.6 | 2.7 | 3.6 | 58.0 |
| IC52599 | Madhya Pradesh | 22.8534 | 79.8434 | 28.0 | 25.0 | 4.5 | 99.0 | 29.5 | 2.86 | 4.3 | 55.0 |
| IC52600 | Madhya Pradesh | 23.5451 | 81.7208 | 43.4 | 27.8 | 4.1 | 49.2 | 44.8 | 2.86 | 5.8 | 76.0 |
| IC16225 | Maharashtra | 19.8891 | 77.9166 | 37.0 | 22.7 | 3.8 | 63.8 | 37.8 | 2.64 | 3.7 | 47.2 |
| IC16236 | Maharashtra | 18.6045 | 76.6814 | 15.5 | 19.0 | 4.0 | 58.0 | 16.2 | 2.6 | 1.96 | 39.7 |
| IC16238 | Maharashtra | 17.9623 | 75.5451 | 59.0 | 30.2 | 4.0 | 97.0 | 60.4 | 2.82 | 9.18 | 55.6 |
| IC16243 | Maharashtra | 19.7409 | 76.8297 | 32.0 | 21.0 | 4.0 | 57.8 | 33.6 | 2.4 | 3.03 | 47.2 |
| IC16248 | Maharashtra | 18.3575 | 75.4957 | 33.0 | 18.0 | 4.0 | 59.6 | 34.3 | 2.6 | 4.68 | 54.6 |
| IC16249 | Maharashtra | 19.6421 | 80.3869 | 47.0 | 18.0 | 4.0 | 47.5 | 48.0 | 3.0 | 9.7 | 69.0 |
| IC16250 | Maharashtra | 20.729 | 79.0529 | 48.0 | 27.0 | 4.0 | 58.0 | 48.7 | 2.9 | 9.93 | 68.5 |
| IC41906 | Maharashtra | 20.482 | 76.4344 | 67.6 | 19.0 | 4.0 | 79.2 | 68.2 | 2.6 | 7.17 | 40.8 |
| IC41910 | Maharashtra | 18.7033 | 77.0273 | 42.2 | 25.4 | 4.0 | 58.6 | 43.2 | 2.74 | 6.7 | 59.0 |
| IC41911 | Maharashtra | 21.0748 | 78.7565 | 31.0 | 22.7 | 3.7 | 54.2 | 31.2 | 2.7 | 4.51 | 54.0 |
| IC41912 | Maharashtra | 20.3831 | 76.5826 | 36.8 | 24.6 | 4.5 | 47.0 | 36.8 | 2.75 | 2.9 | 29.4 |
| IC41978 | Maharashtra | 17.7152 | 73.8159 | 56.3 | 23.8 | 4.0 | 86.0 | 57.3 | 2.7 | 6.99 | 46.0 |
| IC751 | Maharashtra | 18.9504 | 73.6183 | 29.2 | 22.8 | 3.7 | 63.2 | 30.6 | 2.6 | 4.2 | 54.6 |
| IC20156 | Nagaland | 26.1636 | 94.4181 | 38.0 | 22.2 | 4.1 | 57.6 | 39.0 | 2.68 | 2.08 | 53.4 |
| IC16832 | Punjab | 31.4006 | 75.4957 | 54.0 | 20.3 | 4.2 | 73.6 | 54.8 | 2.7 | 3.91 | 43.2 |
| IC31379 | Punjab | 30.2643 | 75.6933 | 44.2 | 19.8 | 3.6 | 47.8 | 45.2 | 2.7 | 2.5 | 27.2 |
| IC14080 | Rajasthan | 27.0529 | 76.5332 | 28.3 | 21.8 | 4.0 | 44.3 | 28.7 | 2.5 | 4.89 | 69.2 |
| IC14106 | Rajasthan | 26.0154 | 75.1993 | 49.8 | 25.4 | 3.8 | 65.0 | 50.6 | 2.9 | 4.75 | 36.4 |
| IC14135 | Rajasthan | 24.2368 | 74.2112 | 54.6 | 29.6 | 4.4 | 72.6 | 55.2 | 2.58 | 4.7 | 43.5 |
| IC14155 | Rajasthan | 24.9779 | 74.0629 | 54.2 | 27.4 | 4.0 | 69.6 | 54.6 | 2.52 | 6.07 | 44.8 |
| IC14174 | Rajasthan | 28.1893 | 74.5557 | 38.8 | 29.0 | 3.4 | 44.0 | 39.2 | 2.92 | 2.09 | 51.0 |
| IC26303 | Rajasthan | 25.5708 | 72.482 | 47.6 | 28.4 | 4.6 | 96.8 | 48.0 | 2.6 | 5.89 | 47.6 |
| IC42965 | Rajasthan | 26.6083 | 74.9522 | 66.4 | 28.6 | 4.0 | 92.2 | 67.2 | 2.8 | 12.19 | 65.6 |
| IC42987 | Rajasthan | 28.5351 | 75.4463 | 69.2 | 33.2 | 4.2 | 151.4 | 70.8 | 2.9 | 11.8 | 59 |
| IC96098 | Uttar Pradesh | 26.0648 | 79.6952 | 41.4 | 22.4 | 3.4 | 69.6 | 42.4 | 2.62 | 2.3 | 70.8 |
| IC96109 | Uttar Pradesh | 27.4482 | 78.8059 | 41.4 | 24.2 | 4.4 | 86.2 | 42.4 | 2.93 | 4.04 | 62.2 |
| IC96113 | Uttar Pradesh | 29.0786 | 78.4601 | 33.2 | 22.9 | 3.9 | 46.4 | 34.4 | 2.6 | 2.28 | 45.6 |




Fig 2. Diversity index for number of capsules per plant in sesame




Fig 4. Diversity index for seed yield in sesame

and Madhya Pradesh are diversity rich pockets for sesame.

Coefficient of variation

High coefficient of variation for plant height (25.0–42.0%) was found in accessions from Madhya Pradesh, followed by a moderate CV% range of 14.0–25.0% in Maharashtra. However, low CV% (up to 8.0%) was recorded on sesame germplasm accessions from other states. Gujarat and Maharashtra recorded low to moderate CV% for the quantitative trait of internode length, which indicated that diversity is less for this trait. Grids with high CV% for the traits, number of leaves plant⁻¹ (29.0–37.0%) and number of flowers plant⁻¹ were stretched in Madhya Pradesh and Maharashtra. Germplasm accessions augmented from Gujarat, Madhya Pradesh and Maharashtra recorded highest CV% (35.0–44.0%) for number of capsules plant⁻¹. Accessions with maximum CV (18.0–23.0%) recorded for seeds capsule⁻¹ originated from Maharashtra, while moderate CV value of 9.0–18.0% was possessed by the germplasm accessions from Madhya Pradesh. Grid map generated for seed weight indicated a lower CV% value (2.0–8.0%) for the accessions studied. This indicated less variability among the accessions. However, maximum CV% value




Fig 3. Coefficient of variation (%) for number of capsules per plant

Madhya Pradesh only. The Shannon diversity index ranged from 0.55–1.0 for germplasm accessions sourced from Gujarat, Madhya Pradesh and Maharashtra states. Among the sesame germplasm sourced from eight Indian states, and based on DIVA-GIS analysis it can be concluded that Gujarat, Madhya Pradesh

(6.0–8.0%) was recorded for Madhya Pradesh collections. The seed yield recorded the highest CV% (48.0–60.0%) among all the plant traits studied.

The present study revealed that diverse germplasm accessions of sesame were dispersed all over the eight states (Gujarat, Himachal Pradesh, Madhya Pradesh, Nagaland, Punjab, Rajasthan and Uttar Pradesh) but region where the diverse genotypes most likely to found are Maharashtra, Madhya Pradesh and Gujarat as high diversity indices and CV% were recorded for all the traits studied. It may be concluded that Maharashtra, Madhya Pradesh and Gujarat states are most suitable for future germplasm collection missions. The present study would be the first of its kind in India by integrating DIVA-GIS for the study of diversity analysis in sesame.

GIS mapping may be effectively used for documentation, diversity analysis, identifying gaps in collection, assessment of loss of diversity, developing new strategies for conservation, and sustainable utilization, particularly in the wake of recent international developments related to food and nutritional security. GIS mapping has been successfully used in assessing biodiversity and in identifying areas of high diversity in *Phaseolus* bean (Jones *et al.* 1997), wild potatoes (Hijmans & David *et al.* 2001), horsegram (Sunil *et al.* 2008), *Jatropha curcas* (Sunil *et al.* 2009), linseed (Sivaraj *et al.* 2009), blackgram (Babu Abraham *et al.* 2010), *Canavalia* fatty acids (Sivaraj *et al.* 2010), medicinal plants (Varaprasad *et al.* 2007), *Piper* (Utpala *et al.* 2006) and agrobiodiversity (Varaprasad *et al.* 2008).

References

- Babu Abraham V, Kamala N, Sivaraj N, Sunil N, Pandravada S R, Vanaja M & Varaprasad, K S 2010 DIVA-GIS approaches for diversity assessment of pod characteristics in black gram (*Vigna mungo* L. Hepper). Cur. Sci. 98: 616–619.
- Hijmans R J, Garret K A, Huaman, Z, Zhang D P, Schreuder M & Bonierbale 2000 Assessing the geographic representatives of genebank collections: the case of Bolivian wild potatoes. Conserv. Biol. 14: 1755–1765.
- Hijmans R J & David M 2001 Geographic distribution of wild potato species. Amer. J. Bot. 88: 2101–2112.
- Jones P G, Beebe S E, Tohme J & Galway N W 1997 The use of geographical information systems in biodiversity exploration and conservation. Biodiv. Conserv. 6: 947–958.
- Sivaraj N, Sunil N, Pandravada S R, Kamala V, Rao B V S K, Prasad R B N, Nayar E R, Joseph John K, Abraham Z & Varaprasad, K S 2010 Fatty acid composition in seeds of Jack bean [*Canavalia ensiformis* (L.) DC] and Sword bean [*Canavalia gladiata* Jacq.] DC] germplasm from South India: A DIVA-GIS analysis. Seed Tech. 32: 46–53.
- Sunil N, Sivaraj N, Pandravada S R, Kamala V, Raghuram Reddy P & Varaprasad K S 2008 Genetic and geographical divergence in horsegram germplasm from Andhra Pradesh, India. Plant Genetic Resources: Characterization and Utilization 7: 84–87.
- Sunil N, Sivaraj N, Anitha K, Babu Abraham, Vinod Kumar E, Sudhir M, Vanaja & Varaprasad K S 2009 Analysis of diversity and distribution of *Jatropha curcas* L. germplasm using Geographic Information System (DIVA-GIS). Genet. Resour. Crop Evol. 56: 115–119.
- Utpala P, George J K, Saji K V, Srinivasan V, Madan M S, Prem Nath Mathur, Parthasarathy V A 2006 Spatial analysis for *Piper* species distribution in India. Pl. Genet. Resour. Newslet. 147: 1–5.
- Varaprasad K S, Sivaraj N, Mohd Ismail & Pareek S K 2007 GIS mapping of selected medicinal plants diversity in the Southeast Coastal Zone for effective collection and conservation. In: Advances in Medicinal Plants (Eds. Janardhan Reddy K, Bir Bahadur, Bhadraiah B & Rao M L N). Universities Press (India) Private Ltd. pp.69–78.
- Varaprasad K S, Sivaraj N, Pandravada S R, Kamala V & Sunil N 2008 GIS mapping of Agrobiodiversity in Andhra Pradesh. Proceedings of Andhra Pradesh Academy of Sciences. Special Issue on Plant wealth of Andhra Pradesh. pp 24–33.